




Product Information Sheet



X877 X-Gluc, Monocyclohexyl Ammonium Salt

Synonym: 5-Bromo-4-chloro-3-indolyl- β -D-glucuronic Acid,
Monocyclohexyl Ammonium Salt

CAS: 114162-64-0

Formula: $C_{20}H_{26}BrClN_2O_7$

Molecular Wt: 521.8

Properties

Form: Powder

Appearance: White Powder

Application: Molecular Biology

Solubility: DMSO or KOH

Typical Working

Concentration:

Storage Temp: -20 to 0° C

Storage Temp of

Stock Solution: -20 to 0° C

Other Notes:

Application Notes

X-Gluc is used in similar fashion as X-Gal. X-Gluc is used to detect β -glucuronidase activity in genetically modified tissue, primarily plant cells. β -glucuronidase is encoded by the *gusA* (*uidA*) gene found in *Escherichia coli*. When present in non-modified cells, β -glucuronidase cleaves X-gluc, leaving glucuronic acid and an intermediate indoxyl. The intermediate indoxyl oxidizes and forms an insoluble blue dimer, 5,5'-dibromo-4,4'-dichloro-indigo, hence it is an ideal GUS activity indicator.

Revised August 2011 PN

PhytoTechnology Laboratories®

P.O. Box 12205 • Shawnee Mission, KS • 66282-2205

Phone: 1-913-341-5343 or 1-888-749-8682 (U.S. Only) Fax: 1-913-341-5442 or 1-888-449-8682 (U.S. Only)

Web Site: www.phytotechlab.com

© 2007 PhytoTechnology Laboratories®